

Drosophila tsukubaensis, a New Species of the *obscura* Group of the Genus *Drosophila* (Diptera, Drosophilidae) from Japan

Hisaki TAKAMORI

Institute of Biological Sciences, University of Tsukuba, Sakura-mura, Ibaraki, 305 Japan

and

Toyohi OKADA

Gôtokuji 2-30-18, Setagaya-ku, Tokyo, 154 Japan.

Synopsis A new species belonging to the *obscura* group of *Drosophila* (*Sophophora*) is described from Japan. The chromosome configuration is also determined.

In Japan have been known to occur 4 species of the *obscura* subgroup of the *obscura* group of *Drosophila*, namely *alpina* BURLA, 1948, *bifasciata* POMINI, 1940, *imaii* MORIWAKI et OKADA, 1967 and *eniwae* TAKADA, BEPPU et TODA, 1979 (MORIWAKI *et al.*, 1952; OHBA, 1954; OKADA, 1956; MORIWAKI *et al.*, 1967; TAKADA *et al.*, 1979).

Recently not a few flies of the 5th member in Japan of this subgroup were collected from Ibaraki, Tochigi and Nagano Prefectures, Honshu, Japan. These flies are concluded to belong to a new species.

This paper describes the new species with note on metaphase chromosome configuration.

Drosophila (Sophophora) tsukubaensis n. sp.

[Japanese name: Tsukuba-shôjôbae] (Fig. 1)

Male and female. Body about 2.5 (1.8-2.8) mm in length, brownish black. Frons, dull black about half as broad as head width and with a few frontal hairs. Eye dark red, with short black pile. Cheek 1/3-1/4 as broad as the greatest diameter of eye. Anterior reclinate orbital about 1/3 as long as proclinate orbital. Antenna black, second joint paler. Arista with 4 upper and 2 lower branches besides a moderate fork. Palpus greyish brown, with 1 or 2 prominent setae. Carina low and broader below. Second orals 1/3 as long as vibrissa. Mesonotum brownish black. Acrostichal hairs in 8 rows. Length distance of dorsocentrals about half the cross distance. Humeral 2, subequal. Sterno-index about 0.5 (0.4-0.7). Legs dark brown, tibiae and tarsi paler. Preapicals on all tibiae, apicals on middle. Proximal




Fig. 1. *Drosophila tsukubaensis* n. sp. A, Periphallallic organs; B, decasternum; C, phallic organs; D, male internal reproductive organs; E, ejaculatory apodeme; F, egg guide; G, female internal reproductive organs. a, Anterior paramere; c, cercus; e, aedeagus; g, epandrium; h, hypandrium; p, posterior paramere; s, surstylus. Scales: broken line, 1.0 mm; solid line, 0.1 mm.

2 tarsal joints of male fore legs provided with sex combs, composed of about 8 (5–10) and 8(6–11) teeth respectively. Wing hyaline, slightly fuscous along costa. C-index about 2.6(2.2–3.1); 4V-index about 2.1(1.8–2.4); 4C-index about 1.1(0.9–1.3); 5X-index about 2.1(1.5–2.6). C1-bristles 2; C3-fringe on basal 1/5–2/5. Halteres yellowish white. Abdominal tergites brownish black. Sternites greyish.

Periphallallic organs (Fig. 1A). Epandrium brownish black, triangular below, lower portion with about 15 hairs, upper portion about 8. Surstylus with a dark brown hook-shaped projection at the upper portion of its caudal margin, a row of about 10(8–11) black teeth at middle, and about 20 brown stout setae distally. Cercus separated from epandrium, dark brown, elliptical and hairy. Decasternum dark brown, small and triangular (Fig. 1B).

Phallic organs (Fig. 1C). Aedeagus pale brown and bifid. Anterior paramere orange brown, curved, pointed at tip and with a row of about 7 sensilla. Posterior paramere orange brown, pointed at tip and bilobed below. Hypandrium with a pair of short submedian spines. Ventral fragma pale brown, fused to hypandrium and semielliptical. PI=0.6.




Fig. 2. Metaphase chromosomes of male larval brain cell of *D. tsukubaensis*.

Egg-guide (Fig. 1F). Lobe pale yellowish brown, rounded at tip, with about 16(13–22) black marginal and about 2(1–3) black discal teeth and a short subterminal hair. Basal isthmus short, broad and yellowish brown.

Internal structures. Mid intestine coiling about 2 times. Rectal papillae about 1.5 times as long as wide. Malpighian tubes with common stalks short and posterior branches ending free. Testis orange, elongate elliptical, apically rounded or slightly pointed and proximally narrowing. Seminal vesicle banana shaped. Accessory gland transparent and folded once (Fig. 1D). Ejaculatory bulb oval, cephalic and caudal margins weakly bilobed. Ejaculatory apodeme with plate rectangular, anterior margin strongly projected and posterior margin angularly concaved (Fig. 1E). Spermatheca hemispherical and black. Parovarium with head oval. Ventral receptacle short and transversely folded about 2 times (Fig. 1G).

Holotype: ♂, Sakura-mura, Ibaraki Pref., 1. VI. 1981, found in banana trap in the campus of the University of Tsukuba (TAKAMORI). Paratypes: 6 ♂, 28 ♀, Sakura-mura, 13. V. —6. VI. 1981; 3 ♂, 3 ♀, Mt. Tsukuba, Ibaraki Pref., 10. VI. 1981; 2 ♀, Meotobuchi, Tochigi Pref., 28. VIII. 1981 (all by TAKAMORI); 5 ♂, 1 ♀, Matsushiro, Nagano Pref., 4. VIII. 1982 (YABUSAKI). Holotype is deposited in the National Science Museum, Tokyo. Paratypes are deposited in the University of Tsukuba.

Relationships. This species undoubtedly belongs to the *obscura* subgroup in dark body color, having 8 rows of acrostichal hairs on mesonotum, low carina, elliptical and orange colored testis, not a few teeth of distal sex comb of the male and short ventral receptacle of the female, yet can be distinguished from any other known species of the subgroup by small and triangular decasternum of the male and exceptionally short basal isthmus of egg-guides of the female.

Among the Japanese species of the *obscura* subgroup, this species is closest to *D. bifasciata*, which was sometimes collected together with this species. This species can be easily distinguished, however, from *D. bifasciata* by paler tarsi.

