

No.

I. OKADA

From the Proceedings of the Linnean Society of New South Wales,
Vol. iii, Part 2, 1927.

NOTES ON AUSTRALIAN DIPTERA. No. x.

By J. R. MALLOCH.

(Communicated by Dr. E. W. Ferguson.)

(Twelve Text-figures.)

[Read 30th March, 1927.]

In this paper I present a revised synoptic key to the species of the genus *Drosophila* Fallén, with descriptions of some new species, a synopsis of the species of the genus *Homoneura* van der Wulp, descriptions and records of some species of the genus *Sapromyza* Fallén, and descriptions of some other acalyptrate Diptera, most of which have been received from Dr. E. W. Ferguson.

Family Drosophilidae.

Genus MYCODROSOPHILA Oldenberg.

This genus is distinguished from *Drosophila* by the extremely short anterior pair of postsutural dorsocentral bristles, which, like the basal pair of scutellar bristles, are almost indistinguishable and by the humped up thoracic dorsum. All the known species are very similar in colour, glossy black on dorsum, with yellow abdominal markings and pale yellow on venter and sides of thorax. The flies occur commonly on the undersides of fungi in which the larvae feed.

I have before me a series of specimens belonging to a species which is evidently undescribed and one injured specimen which may possibly belong to another species.

MYCODROSOPHILA ARGENTIFRONS n. sp.

Male and female.—Head black, yellow on centre of frons from ocelli to anterior margin and on lower half of occiput, the face brownish, bases of antennae yellowish; orbits shining, rest of frons dull, when seen from in front densely silvery; proboscis yellow below, fuscous above; palpi fuscous. Thorax glossy black on dorsum, with a variable amount of yellow on anterior margin, sometimes in the form of two rudimentary vittae; scutellum when seen from in front brownish or yellowish dusted apically; pleura pale yellow, black on upper margin; postnotum black. Abdomen pale yellow, with a black fascia across hind margin of each tergite which is carried forward on the median line on tergites 3 to 5 and connects with a similar fascia on anterior margin leaving two yellow spots on each, the lateral margins of tergites wholly black. Legs pale yellow. Wings hyaline, with a deep black mark on costa before apex of first vein and a faint cloud extending from it over cell to second vein. Halteres black.

Head a little broader than thorax; frons at vertex nearly twice as wide as length at centre; all bristles, except the anterior reclinate one on each orbit, long; eyes almost bare; facial carina widened and flattened below; clypeus convex. Thorax quite conspicuously convex, the dorsal surface rather densely fine haired. Third costal division over half as long as second.

Length, 2.3 mm.

Type male, allotype, and 5 paratypes, Coramba, N.S.W., 15.2.25.

Genus *LEUCOPHENGA* Mik.
LEUCOPHENGA MINUTA n. sp.

Male.—Head testaceous, occiput and face, except sides, infuscated, frons slightly darkened above, but all these parts with dense white dusting so that in some positions they appear pale; palpi and antennae testaceous; cephalic bristles, except on vibrissal angles, yellow. Thorax testaceous yellow, densely white dusted, almost silvery when seen from certain angles, with evidences of a rufous vitta along each side of dorsum; a brownish vitta along centre of pleura; scutellum testaceous yellow; postnotum brown. Abdomen brownish, when seen from in front with quite dense silvery dusting on dorsum. Thoracic and abdominal hairs and bristles luteous. Legs testaceous yellow. Wings hyaline, a narrow fuscous cloud extending from apex of posterior basal cell obliquely to apex of first vein. Halteres missing in type.

Frons about twice as long as wide, parallel-sided, all bristles well developed; face slightly carinate on upper half; antennae normal; cheek linear; vibrissae short; palpi slender. Both sternopleurals long. Legs slender, no outstanding setulae on mid tibiae. First posterior cell of wing not narrowed apically; third costal section over half as long as second.

Length, 2 mm.

Type, Cairns District, Queensland (Dodd).

Distinguished from other Australian species known to me by the uniformly silvery white dusted dorsum, markings of the wings, and its small size.

Genus *DROSOPHILA* Fallén.

As I have described a number of species of this genus since I published my key in the first paper of this series (These PROCEEDINGS, 1923, p. 615) I deem it expedient to present now an enlarged synopsis which includes 22 species, all at present known to me as occurring in Australia. I have found it necessary to draw up this key for my own convenience in identifying material and hope that it may prove as useful to other students of the group as it has to me for that purpose. There must be many Australian species of the genus unknown to me. Most of those herein included are similar to the general run of species of the genus throughout its range, but some, and especially *nicholsoni*, are aberrant, though not entitled, in my opinion, to subgeneric segregation. I have not attempted to figure the genitalia of either sex, some of which possess striking specific features, nor have I attempted to determine the structure of the eggs of the species. Some of the eggs are furnished with filaments at one end which vary in form and in number with the species and they may sometimes be as readily distinguished as any of the other stages, and in some closely allied species even more readily than the adults.

Key to the Species.

1. Wings with conspicuous fuscous markings in addition to those over the cross veins or at apex of first costal division 2
- Wings without fuscous markings except sometimes faint clouds over the cross veins or at apex of first costal division 3
2. Wing with a large brownish or fuscous spot at apex of second vein, the dark cloud at tip consisting of a brownish suffusion along the apices of third and fourth veins, more or less coalescent in first posterior cell; mesonotum dark brown, with three linear yellow vittae, the median one not reaching anterior margin; scutellum yellow in centre, dark brown on sides of disc; pleura whitish-yellow, contrasting sharply with the dark brown mesonotum *mycetophaga* Malloch.

- Wing without a dark spot at apex of second vein, the dark cloud at tip consisting of a broad curved brown patch which extends from middle of third section of costa to just over third vein and over disc of wing to beyond fourth vein, but leaves a hyaline spot in apex of first posterior cell; mesonotum brown, with two poorly defined paler vittae which are carried over lateral margins of scutellum, the centre of latter brown; pleura not noticeably paler than mesonotum *polyptori* Malloch.
3. Fore femora with a comb-like series of microscopic setulae on apical third or more of anteroventral surface 4
Fore femora without a comb-like series of microscopic setulae as above 5
4. Fore femora with short closely placed setulae on more than the apical half of posteroventral surfaces, the longest one, at apex, not longer than the femoral diameter; third section of costa not less than one-third as long as second; facial carina narrow *setifemur* Malloch.
Fore femora with four or five widely spaced bristles on the entire length of each posteroventral surface, the longest one, at middle, as long as or longer than the femoral diameter; third section of costa about one-fourth as long as second; facial carina much broadened below *immigrans* Sturtevant.
5. Thoracic dorsum with a minute but conspicuous dark brown or fuscous dot at base of each hair and bristle and sometimes with additional brown or fuscous dots on parts causing aggregations which assume the appearance of larger irregular spots, the ground colour of thoracic dorsum never testaceous yellow 6
Thoracic dorsum with or without dark or pale vittae, not copiously marked with piliferous spots or dots, if faintly marked with piliferous dots then the ground colour of thorax is testaceous yellow 9
6. Third costal division of wing almost as long as second; outer cross vein about one-third as long as apical section of fifth vein *poecilothorax* Malloch.
Third costal section of wing not over one-third as long as second; outer cross vein much more than one-third as long as apical section of fifth vein 7
7. Facial carina practically absent except between bases of antennae, where it is sharp and linear; eyes almost bare *obsoleta* Malloch.
Facial carina conspicuous below, where it is rather broad and usually longitudinally sulcate; eyes with dense stiff microscopic hairs 8
8. At least the first three tergites of abdomen with a yellowish spot on each side on the part that is incurved on venter *repleta* Wollaston.
None of the abdominal tergites with yellowish spots as above *hydei* Sturtevant.
9. All hairs and bristles on insect luteous; thorax not vittate *flavohirta* Malloch.
All hairs and bristles fuscous or blackish 10
10. Sides of face, frontal orbits, and two narrow submedian vittae on mesonotum which extend over its entire length and on to sides of scutellum, densely white dusted, the whole forming two continuous white lines on a black ground, and very conspicuous; sternopleurals 2 *albostrigata* Malloch.
Thoracic dorsum with or without dark vittae, not white vittate 11
11. Thoracic dorsum distinctly vittate 12
Thoracic dorsum entirely without vittae 16
12. Thoracic dorsum with five dark vittae, the intervening spaces pale grey dusted; wing with a deep black spot and two fine bristles on costa before apex of first vein; third costal section almost as long as second *nigrovittata* Malloch.
Thoracic dorsum with four or six dark vittae, the intervening spaces inconspicuously pale dusted 13
13. Arista with but one long hair above at base, no long hairs below; face without a carina on lower portion; third costal section not over one-fourth as long as second, and shorter than penultimate section of fourth vein; sternopleurals 3 *nicholsoni*, n. sp.
Arista with at least three long hairs above and one or two below 14
14. Palpi fuscous; face with a vestigial carina only on upper part, not carinate below; pleura with two dark vittae; third costal division over one-third as long as second and distinctly longer than penultimate section of fourth vein; sternopleurals 2 *buscki* Coquillett.
Palpi testaceous yellow; face with a very conspicuous carina which is continued below middle, its lower portion broadened 15

15. Submedian thoracic vittae narrowed at posterior extremities, but continued to the hind margin of mesonotum, no dark central mark in front of scutellum; the two central series of hairs located on the inner margins of the submedian dark vittae *lativittata* Malloch.
 Submedian vittae not reaching posterior margin of mesonotum; a dark mark on centre of hind margin of latter; the two central series of hairs entirely within the pale central vitta, the next two series on edges of the dark submedian vittae *enigma* n. sp.
16. Facial carina not developed except weakly on upper part between bases of antennae 17
 Facial carina well developed and broadened on lower part of face 18
17. Thorax dull fuscous *fuscithorax* Malloch.
 Thorax testaceous yellow, sometimes with faint dark dots at bases of the bristles and hairs on dorsum *mornata* Malloch.
18. Thorax more or less shining black 19
 Thorax shining testaceous yellow 21
19. Third costal section distinctly more than half as long as second and longer than penultimate section of fourth vein (3:2); frons reddish or yellowish brown except on orbits and triangle, which are shining black, the anterior reclinate orbital bristle laterad of the proclinate one *sydneyensis*, n. sp.
 Third costal section about one-third as long as second and about as long as, or very little longer than, penultimate section of fourth vein 20
20. Frons dark brown, the orbits and triangle dull black; anterior reclinate orbital bristle well above the proclinate one; thorax shining, but not glossy *subnitida* n. sp.
 Frons black, the orbits and triangle shining, anterior reclinate orbital bristle but little above level of the proclinate one; thorax glossy *nitidithorax*, n. sp.
21. Fore metatarsus of male with a comb of short black bristles on apical half, the second segment without a comb *ampelophila* Loew.
 Fore metatarsus and second segment each with a comb of black bristles from base to apex in male *serrata*, n. sp.

[*Drosophila brunneipennis* Malloch described in These PROCEEDINGS, 1923, p. 617, has inadvertently been omitted. In the previous key *loc. cit.* p. 616 it is placed next to *D. melanogaster* Meigen = *D. ampelophila* Loew.—Ed.]

Scaptomyza

✓ *DROSOPHILA NICHOLSONI*, n. sp.

Male and female.—Testaceous yellow, shining. Frons opaque brownish, the orbits, sides of triangle, and anterior margin testaceous, not shining; third antennal segment largely fuscous; palpi and proboscis testaceous. Dorsum of thorax slightly brownish and greyish dusted with six brown vittae, the submedian pair of moderate width and complete, the intermediate pair much narrower and connected with the submedian pair behind suture, the sublateral pair broad, rather indistinct and broken at suture; pleura and metanotum largely brown; scutellum brown in centre, narrowly yellow on sides. Abdomen shining, apical tergites slightly darkened. Legs testaceous yellow. Wings greyish hyaline, veins not clouded. Halteres yellow.

Anterior reclinate bristle about one-third as long as posterior one and almost in transverse line with the proclinate one; postvertical bristles long interfrontalia with a few hairs anteriorly; facial carina distinguishable only on upper part; arista with one long hair on upper side at base and some short pubescence beyond, which is only visible under a high magnification; vibrissa single. Thorax with two humerals, about eight series of intradorsocentral setulae, the submedian two series on margins of the pale central vitta, and 3 sternopleurals. Legs normal. Second section of costa about four times as long as third, the latter not longer than penultimate section of fourth vein, ultimate section of fifth vein subequal to penultimate section of fourth and about twice as long as outer cross vein.

Length, 2.5-3 mm.

Type male, allotype, and 2 paratypes, Perth, W.A., 15.11.1924 (Nicholson).

This species is readily distinguished from any other known species in the genus either from Australia or elsewhere by the presence of but one long hair on the upper side of the arista. This character was used as the distinguishing feature of the American genus *Cladochaeta* by Coquillett, but the present species differs from that genus in many characters and is so obviously merely an aberrant *Drosophila* that I retain it in this genus, while noting that it apparently weakens the claim of *Cladochaeta* to generic recognition.

Scaptomyza

✓ DROSOPHILA NITIDITHORAX, n. sp.

Male.—Deep black, dorsum of thorax and of abdomen almost glossy. Face whitish dusted and, like the cheeks, partly brownish yellow; frons except the orbits and triangle velvety black; antennae black; palpi and proboscis dusky testaceous yellow. Thorax without markings, the pleura not so conspicuously shining as dorsum. Abdomen slightly yellow at base. Legs dusky testaceous yellow, femora almost entirely fuscous. Wings greyish hyaline, veins unclouded. Halteres brownish yellow.

Frontal bristles strong, anterior reclinate orbital half as long as posterior one and slightly but distinctly above level of the proclinate one; interfrontalia with quite dense short black setulose hairs; facial carina nose-like, rounded above on the lower part; rays of arista about 4+2; vibrissae strong. Intradorsocentral setulae in 8-10 series; sternopleurals 3. Legs normal. Third costal section not longer than penultimate section of fourth vein and about one-third as long as second section; outer cross vein at about 1.5 its own length from apex of fifth vein.

Length, 2.5 mm.

Type and paratype, Perth, W.A., 15.11.1924 (Nicholson).

A robust black species which is most closely related to *subnitida* described below.

Scaptomyza

✓ DROSOPHILA SUBNITIDA, n. sp.

Female.—Distinguished from the preceding species by the brown colour of the frons, the opaque frontal orbits and less shining dorsum of thorax.

There are no outstanding structural distinctions, but the anterior reclinate bristle is larger and farther from the proclinate one, the third section of costa is a little longer and the insect is less robust.

Length, 2 mm.

Type, Sydney, N.S.W., 6.1.25.

This species has somewhat the appearance of *fuscithorax* Malloch, but the latter has no carina on lower part of face and differs in other respects.

Scaptomyza

✓ DROSOPHILA SYDNEYENSIS, n. sp.

Female.—Black, dorsum of thorax and abdomen almost glossy. Frons brownish red, orbits and triangle shining black; face yellowish on sides, slightly white dusted; cheeks yellowish; antennae yellowish or rufous, third segment fuscous; palpi and proboscis yellowish. Thorax as in *nitidithorax*, but the pleura as conspicuously shining as mesonotum. Abdomen with bases of basal two or three tergites yellowish. Legs as in *nitidithorax*. Wings hyaline. Halteres yellow.

Frons with a few hairs in front; anterior reclinate orbital bristle short, in transverse line with proclinate one; facial carina well developed and nose-like.

Thorax as in *subnitida*. Legs normal. Wing venation differing from that of *nitidithorax* and *subnitida* in having the third costal division well over half as long as second, usually about, or even full, three-fourths as long as it, and twice as long as penultimate section of fourth vein, the latter subequal to ultimate section of fifth, and about twice as long as outer cross vein.

Length, 2 mm.

Type and three paratypes, Sydney, N.S.W., 3 and 5.1.1925, and 2.4.1925.

A less robust species than *nitidithorax*.

Scaptomyza

✓ *DROSOPHILA ENIGMA*, n. sp.

Male and female.—Very similar to *lativittata* Malloch in colour and general structure. Differs from it in being paler, the ground colour being testaceous yellow, the dorsum of thorax greyish dusted, and with paler vittae which are narrower, the submedian pair separated by a wider space which covers four instead of two series of the short hairs, and the other characters of the markings as stated in the key. The abdominal markings consist of a dark brown fascia on hind margin of each tergite which is narrowed or interrupted in centre, widened each side of median line and again at the lateral curvature, the fasciae rather indistinct on the lateral incurved portions of the tergites. As in *lativittata* the outer cross vein is faintly clouded and the penultimate section of fourth vein is about half as long as ultimate.

Length, 3 mm.

Type male and allotype, Sydney, N.S.W., 22.7.23, and 23.9.24. Paratypes, two females, Toronto, N.S.W.

✓ *DROSOPHILA SERRATA*, n. sp. (Text-figure 1.)

Male.—Shining fulvous yellow, very similar to *ampelophila* Loew which it closely resembles in many respects. The type specimen is rather immature, but the thorax is not vittate and the abdomen has a faint dark uninterrupted apical fascia on each tergite. The legs are yellow and the wings yellowish hyaline with unclouded veins. Halteres yellow. Bristles and hairs fuscous.

Anterior reclinate orbital bristle about one-third as long as posterior one and distinctly above the level of proclinate one; eyes quite densely stiff-haired; face distinctly carinate. Thorax with six series of intradorsocentral hairs; prescutellar acrostichals not differentiated; both humerals strong; only two sternopleurals well developed. Two basal segments of fore tarsus with a comb-like series of short stiff



Text-fig. 1. *Drosophila serrata*, two basal segments of fore tarsi of male from in front.

black bristles on anterior side, the comb on basal segment bipartite (Fig. 1). Third section of costa about half as long as second and a little less than twice as long as penultimate section of fourth vein, the latter about one-third as long as ultimate section and subequal to ultimate section of fifth vein; outer cross vein at about twice its own length from apex of fifth vein.

Length, 1.75 mm.

Type, Eidsvold, Queensland, 2.4.24 (Bancroft).

There are several described species of the genus with the two basal segments of fore tarsi armed with comb-like bristles, but none of these have the armature as in this species so far as I am aware. I described one, *biarmipes*, from India, but in it the combs are confined to the apical parts of each segment and do not extend along the whole length of the anterior surfaces as here.

Genus GITONIDES Knab.

The species described below falls most readily into *Gitonides*, but it differs from the genotype in having the frons much narrower anteriorly and the first posterior cell of the wing quite noticeably narrowed apically.

Annotata (Erima) fasciata

GITONIDES CONVERGENS, n. sp.

telli Bock, 1982
Annotata & *Erima* *Suzuki*.
 89: 23

Head brownish testaceous; upper half of occiput fuscous, lower half white dusted; upper extremities of frontal orbits, ocellar spot and clypeus, fuscous; face slightly white dusted; palpi brownish, paler at apices. Thorax brownish testaceous, dorsum when seen from in front with a broad brownish central vitta which is faint in front of the suture, where it is sometimes divided centrally and which is divided between suture and hind margin, a broad branch curving to each side and between the two pairs of dorsocentrals, and a narrower central one continuing to hind margin, the disc laterad of the anterior and posterior portions of vitta white dusted, between these pale markings and lateral margins there is a broad brownish vitta; humeral angles testaceous; scutellum when seen from in front with a brown central line which broadens out and covers apex, a fainter brown mark on each basal angle and a less distinct dark mark on disc each side of the dark central line, the latter surrounded by whitish dusting. Abdomen testaceous, all tergites except the basal two largely or entirely black. Legs testaceous, femora browned, tibiae dark at apices, the mid pair most conspicuously, so. Wings hyaline. Halteres yellow.

Frons at vertex about one-third of the head width, much narrowed anteriorly, at front margin not half as wide as long in centre; proclinate orbital bristle well above middle of frons, the anterior reclinate bristle quite small and about midway between the others; arista bare; face slightly carinate; antennae normal, inserted at middle of profile. Thorax with two pairs of postsutural dorsocentrals, the anterior pair short, about eight series of intradorsocentral setulae, one humeral and two sternopleurals; scutellum convex. Abdomen broad and short. Legs stout. normal. Inner cross vein at two-fifths from apex of discal cell; outer cross vein at less than its own length from apex of fifth vein; ultimate section of fourth vein over three times as long as penultimate section; first posterior cell quite noticeably narrowed apically.

Length, 3 mm.

Type, Eidsvold, Queensland, 1924 (Bancroft).

Family Agromyzidae.*

Subfamily MILICHIINAE.

DESMOMETOPA VARIPALPIS, n. sp.

Female.—Head black, whitish-grey dusted, with the usual opaque black M-shaped frontal marking; cheeks yellowish below; palpi testaceous, with conspicuous irregular black spotting. Thorax black, slightly shining, evenly greyish dusted and without vittae. Abdomen greasy in type, black, and probably less dusted than thorax. Legs black, tarsi yellowish, darker at apices. Wings hyaline. Knobs of halteres yellow.