

Makino, Sajiro, and Kanehisa,  
Takeharu A preliminary survey  
of the geographical distribution  
of *Drosophila* in Hokkaido.  
DIS 25:110-111 (1951)  
Central, East, and North.

The species of *Drosophila* so far col-  
lected from Hokkaido and their distri-  
bution are preliminarily listed below.  
For description, the island of Hokkaido  
was divided into four regions-- South

	South	Central	East	North
<i>Amiota</i> sp.		+		
<i>Scaptomyza</i> sp.	+	+		+
<i>Hirtodrosophila</i> sp.		+	+	+
<i>H. cinerea</i> group	+			
<i>melanogaster</i>		+		
<i>auraria</i>	+	+	+	+
<i>suzukii</i>	+	+	+	+
<i>obscura</i> group		+	+	
<i>transversa</i>	+	+	+	+
<i>nigromaculata</i>	+	+	+	+

	<u>South</u>	<u>Central</u>	<u>East</u>	<u>North</u>
virilis	+	+	+	+
testacea	+	+	+	+
funebri	+	+	+	+
histrion		+	+	
immigrans		+	+	
rufa		+	+	
repleta group			+	
melanica group			+	
macularis group			+	
robusta group	+	+		
sordidula		+	+	+
coracina		+	+	
busckii	+	+	+	
Mycodrosophila sp.			+	
grandis (?)			+	